

A STRUCTURE THEOREM AND THE GRADED BETTI NUMBERS FOR ALMOST COMPLETE INTERSECTIONS

ALFIO RAGUSA AND GIUSEPPE ZAPPALÀ

ABSTRACT. We provide a structure theorem for all almost complete intersection ideals of depth three in any Noetherian local ring. In particular, we find that the minimal generators are the pfaffians of suitable submatrices of an alternating matrix. From the graded version of the previous result, we characterize the graded Betti numbers of all 3-codimensional almost complete intersection schemes of $\mathbb{P}^r$.

INTRODUCTION

It is well known that in any regular local ring R every ideal I for which the number of elements in a minimal set of generators is equal to its height is generated by a regular sequence, in particular it is a perfect ideal (in the sense that $\text{hd } R/I = \text{depth } I$). This simple fact has an important consequence for studying projective schemes X of $\mathbb{P}_k^r$, since this class of ideals characterizes complete intersection schemes, i.e. schemes X of codimension c whose defining (saturated) ideal I_X in $R = k[x_0, x_1, \dots, x_r]$ is generated by c elements, hence schemes which can be described with the minimum number of equations. Thus, complete intersection schemes are necessarily generated by a regular sequence and consequently are arithmetically Cohen Macaulay (aCM). In particular, from this one has a very simple description of a minimal free resolutions of R/I_X (Koszul resolution) and consequently of the graded Betti numbers. The question becomes immediately more complicated when one admits that the ideal I_X has one generator more than its depth. In some part of the literature aCM schemes X of $\mathbb{P}_k^r$, of codimension c whose defining ideal I_X is minimally generated by $c + 1$ elements are called *almost complete intersection* schemes. Similarly, in any unitary commutative local ring A a perfect ideal I for which $\nu(I) = \text{depth } I + 1$ is said to be an *almost complete intersection* ideal. Some discussion about almost complete intersection schemes can be found on the J. Migliore book's [Mi]. Very little is known about almost complete intersection schemes (or ideals), for instance some result for “generic” almost complete intersections can be found on the article of J. Migliore and R. Mir Roig [MM]. The very first observation which makes these ideals (or these schemes) nice to study is the fact that every almost complete intersection is directly linked in a complete intersection to a Gorenstein ideal (or aG scheme). Indeed, if $I_Q \subset R$ is the defining ideal of an almost complete intersection of codimension c and $I_Z \subseteq I_Q$ is generated by c minimal generators of I_Q which form a regular

2000 *Mathematics Subject Classification.* 13 D 40, 13 H 10.

Key words and phrases. Almost complete intersections, Gorenstein rings, pfaffians, Betti numbers.

sequence then $I_G := I_Z : I_Q$ is the defining ideal of an aG scheme. By liaison theory (see for instance [PS]) we have also $I_Q = I_Z : I_G$. Now, using this fact and the very nice structure theorem for aG schemes of codimension 3 (Buchsbaum-Eisenbud [BE]), in this paper we will give a structure theorem for all almost complete intersection of depth 3 (see Theorem 2.3). This result generalizes an analogous one of S. Seo which, in [Se] Theorem 2.4, gives a similar characterization for those almost complete intersections Q such that the aG scheme defined by $I_G := I_Z : I_Q$ has the three minimal generators of smallest degree performing a regular sequence. On the other hand, by Diesel paper (see [Di]), one can characterize all graded Betti numbers for 3-codimensional aG schemes. Then, using this characterization and the mentioned structure theorem, we obtain the main result of this paper, i.e a characterization of all graded Betti numbers for 3-codimensional almost complete intersections (Theorem 3.6). Thus, this result provides a new large class of 3-codimensional projective schemes, besides Gorenstein schemes, for which one is able to give a complete description of the graded Betti numbers.

1. NOTATION AND PRELIMINARIES

Let R be a Noetherian local ring. A perfect ideal I of R is called an *almost complete intersection* ideal if $\nu(I) = \text{depth } I + 1$ (here $\nu(I)$ denotes the number of minimal generators of I). Of course, in case R is Cohen-Macaulay an ideal I is an almost complete intersection when $\nu(I) = \text{ht}(I) + 1$, where $\text{ht}(I)$ is its height or codimension. Analogously, let k be an algebraically closed field and $X \subset \mathbb{P}_k^r$ be a closed subscheme of codimension c . Let $I_X \subset R = k[x_0, x_1, \dots, x_r]$ be the saturated homogeneous ideal defining X . Then $X \subset \mathbb{P}_k^r$ is said an *almost complete intersection* scheme when its defining ideal I_X is perfect and minimally generated by $c + 1$ elements. Our first observation is that every almost complete intersection is directly linked in a complete intersection to an arithmetically Gorenstein (aG) scheme. Indeed, if $I_Q \subset R$ is the defining ideal of an almost complete intersection of height c and $I_Z \subseteq I_Q$ is generated by c minimal generators of I_Q which form a regular sequence then $I_G := I_Z : I_Q$ is the defining ideal of an aG scheme. By liaison theory (see [PS]) for a complete discussion on this argument) we have also $I_Q = I_Z : I_G$.

Therefore many properties of almost complete intersections can be deduced by properties of aG schemes. Since in case of codimension $c = 3$ we have the Eisenbud Buchsbaum structure theorem for Gorenstein algebras (see [BE]), in this paper we will produce an analogous structure theorem for all almost complete intersections of depth 3. We will give also a graded version of this structure theorem. Moreover, since again in codimension $c = 3$ by Diesel paper (see [Di]) all graded Betti numbers for Gorenstein graded algebras are characterized, we will describe all possible Betti sequences for almost complete intersections of depth 3. For all of this we need to fix some terminology and to remind some known facts about free resolutions, in particular, for Gorenstein algebras and about pfaffians of alternating matrices. Moreover, since the Betti sequences are simply sequences of finite multisets of positive integers, we need also to fix terminology and basic facts on multisets.

We start just with basic notation and properties of multisets.

Let A be a set. A *multiset* on A is a function $M : A \rightarrow \mathbb{Z}^+$; for every $a \in A$ the integer $M(a)$ will be called the *multiplicity* of a in M and we will denote it also with $\mu_M(a)$. When M is a multiset the domain of M is called the *support* of M and we denote it by $\text{Supp } M$. Whenever $b \notin \text{Supp } M$ we will set $\mu_M(b) = 0$.

Sometimes, for simplicity, we will denote the multiset M with a finite support by $M = \{\{m_1, m_2, \dots, m_t\}\}$, where in the list each element of the support of M appears as many times as its multiplicity in M , and sometimes we will use the notation $M = \{a_1^{(\mu_M(a_1))}, a_2^{(\mu_M(a_2))}, \dots, a_r^{(\mu_M(a_r))}\}$, where a_i are all the elements in $\text{Supp } M$. We define $|M| = \sum_{a \in \text{Supp } M} \mu_M(a)$. Of course, if M is a multiset with $\mu_M(a) = 1$ for all elements in $\text{Supp } M$ we can identify the multiset M with its support.

Now we will remind the definitions concerning the main operations on multisets. If M and N are two multisets we will denote

$$\begin{aligned} M \cap N : \text{Supp } M \cap \text{Supp } N &\rightarrow \mathbb{Z}^+, \quad \text{where } (M \cap N)(y) = \min\{\mu_M(y), \mu_N(y)\}, \\ M \cup N : \text{Supp } M \cup \text{Supp } N &\rightarrow \mathbb{Z}^+, \quad \text{where } (M \cup N)(y) = \max\{\mu_M(y), \mu_N(y)\}, \\ M \sqcup N : \text{Supp } M \cup \text{Supp } N &\rightarrow \mathbb{Z}^+, \quad \text{where } (M \sqcup N)(y) = \mu_M(y) + \mu_N(y), \end{aligned}$$

if we set $\text{Supp}(M \setminus N) = \{x \in \text{Supp } M \mid \mu_M(x) > \mu_N(x)\}$ then

$$M \setminus N : \text{Supp}(M \setminus N) \rightarrow \mathbb{Z}^+, \quad \text{where } (M \setminus N)(y) = \mu_M(y) - \mu_N(y);$$

moreover, we will say that M is a submultiset of N and we will write $M \subseteq N$ whenever $\text{Supp } M \subseteq \text{Supp } N$ and for every $y \in \text{Supp } M$ $\mu_M(y) \leq \mu_N(y)$. Of course, when $M \subseteq N$ then $N = M \sqcup (N \setminus M)$.

In the sequel when M is a finite multiset of integers we define $\|M\| = \sum_{y \in \text{Supp } M} \mu_M(y)y$. Furthermore, if n is also an integer, we define the multiset $n \pm M$ by

$$n \pm M : n \pm \text{Supp } M \rightarrow \mathbb{Z}^+, \quad \text{where } (n \pm M)(n \pm y) = \mu_M(y).$$

Proposition 1.1. *If M is a multiset of integers, n is an integer and $H := M \cap (n - M)$, then for every $x \in \text{Supp } H$ $\mu_H(x) = \mu_H(n - x)$.*

Proof. Of course, for every $x \in \text{Supp } M$ $\mu_M(x) = \mu_{n-M}(n - x)$. Then for every $x \in \text{Supp } H$ we have

$$\begin{aligned} \mu_H(x) &= \min\{\mu_M(x), \mu_{n-M}(x)\} = \\ &= \min\{\mu_{n-M}(n - x), \mu_M(n - x)\} = \mu_H(n - x). \end{aligned}$$

□

Let now $R = k[x_0, \dots, x_r]$ and I_X the defining (saturated) ideal of a projective scheme $X \subset \mathbb{P}_k^r$. We recall that the standard graded k -algebra R/I_X admits a graded minimal free resolution of the following type

$$0 \rightarrow \bigoplus_j R(-j)^{\beta_{pj}} \rightarrow \dots \rightarrow \bigoplus_j R(-j)^{\beta_{0j}} \rightarrow R \rightarrow R/I_X \rightarrow 0$$

which, if we restrict ourselves to the $\beta_{ij} \neq 0$, can be written as

$$0 \rightarrow \bigoplus_{h \in \text{Supp } M_p} R(-h)^{\mu_p(h)} \rightarrow \dots \rightarrow \bigoplus_{h \in \text{Supp } M_1} R(-h)^{\mu_1(h)} \rightarrow R \rightarrow R/I_X \rightarrow 0$$

where each M_i is a multiset of positive integers and $\mu_i(h) := \mu_{M_i}(h)$. We will call $(M_1, \dots, M_p)$ the *Betti sequence* of X or also of R/I_X , and we will denote it by β_X or β_{R/I_X} .

For simplicity, from now on for every multiset of positive integers M we will set

$$\bigoplus_{h \in M} R(-h) := \bigoplus_{h \in \text{Supp } M} R(-h)^{\mu_M(h)}$$

hence, the resolution can be written

$$0 \rightarrow \bigoplus_{h \in M_p} R(-h) \rightarrow \dots \rightarrow \bigoplus_{h \in M_1} R(-h) \rightarrow R \rightarrow R/I_X \rightarrow 0$$

Now, since Gorenstein algebras of depth 3 can be described by alternating matrices, we establish some terminology for such matrices.

Let R be any commutative ring.

If $i \neq j$ are positive integers we set $\langle i, j \rangle$ the following integer

$$\langle i, j \rangle = \begin{cases} i + j + 1 & \text{if } i < j \\ i + j & \text{if } i > j \end{cases}$$

If $M = (a_{ij})$ is an alternating matrix of size m with entries in R we will denote by $M_{\widehat{ij}}$ the alternating matrix obtained from M by deleting both rows and columns i and j . With this terminology when m is even one can easily verify that the pfaffian of M (for definitions and basic facts on pfaffians see for instance [IK] Appendix B) can be computed, for every $i = 1, \dots, m$, by

$$\text{pf } M = \sum_j (-1)^{\langle i, j \rangle} a_{ij} \text{ pf } M_{\widehat{ij}}. \quad (1)$$

Moreover, if M is an alternating matrix of size m , with m even, we will denote by $\overline{M} = (\overline{a}_{ij})$, where $\overline{a}_{ij} = (-1)^{\langle i, j \rangle} \text{pf } M_{\widehat{ij}}$, the pfaffian adjoint of M , which is clearly an alternating matrix.

Remark 1.2. If

$$M = \left(\begin{array}{cc|c} 0 & a & B \\ -a & 0 & \\ \hline - & - & - \\ -{}^t B & & C \end{array} \right)$$

is an even alternating matrix, by repeating the previous formula for $i = 1, 2$, one can show that its pfaffian can be computed as

$$\text{pf } M = a \text{ pf } C + \text{pf}(B \overline{C} {}^t B)$$

where $\overline{C}$ is the pfaffian adjoint of C as defined above.

Let $\psi : F^\vee \rightarrow F$ be an alternating map with F a free R -module of rank even m . If B is a basis for F and $B^\vee$ the dual basis for $F^\vee$, if M is the matrix associated to ψ with respect to such bases, we will denote by $\overline{\psi} : F \rightarrow F^\vee$ the map whose associated matrix with respect to the previous bases is the pfaffian adjoint of M , i.e. $M(\overline{\psi}) = \overline{M}$. Note that $\overline{\psi}\psi = (\text{pf } \psi) \text{id}_{F^\vee}$ and $\psi\overline{\psi} = (\text{pf } \psi) \text{id}_F$. Moreover we will write $\text{Pf}_s(\psi)$ for the ideal generated by the pfaffians of ψ of order s .

Furthermore, with the same notation, when the rank m is odd and $\{p_1, \dots, p_m\}$ are the m submaximal pfaffians of the matrix $M = M(\psi)$ (with respect to

fixed bases B and $B^\vee$), if $p \in I = (p_1, \dots, p_m)$, then there exists an alternating map $\tilde{\psi} : G^\vee \rightarrow G$, where G is a free module of rank $m + 2$, such that the alternating matrix $\tilde{M} = M(\tilde{\psi})$ (with respect to suitable bases $\tilde{B}$ and $\tilde{B}^\vee$) has the $m + 2$ submaximal pfaffians $\{p_1, \dots, p_m, p, 0\}$. Indeed, if $p = a_1 p_1 + \dots + a_m p_m$, it is enough to take $G = F \oplus R^2$ and as $\tilde{\psi} : G^\vee \rightarrow G$, the alternating map which, respect to given bases $\tilde{B}$ and $\tilde{B}^\vee$, is defined by the alternating matrix

$$\tilde{M} = \left(\begin{array}{ccc|cc} & & & 0 & a_1 \\ & M & & \vdots & \vdots \\ & & & 0 & a_m \\ \hline & & & 0 & -1 \\ 0 & \cdots & 0 & 0 & 0 \\ \hline -a_1 & \cdots & -a_m & 1 & 0 \end{array} \right)$$

To complete the assertion it is enough to compute the submaximal pfaffians of $\tilde{M}$, using the above formula (1) with respect to the $(m + 1)$ -th column for all pfaffians except for the $(m + 1)$ -th pfaffian for which we use formula (1) with respect to the last column. Applying the previous observation we get the following

Lemma 1.3. *Let R be a noetherian local ring and I_G a Gorenstein ideal of depth 3 in R and $p_1, p_2, p_3 \in I_G$. Then there exists an alternating map $\varphi : H^\vee \rightarrow H$ of odd rank m such that $\text{Pf}_{m-1}(\varphi) = I_G$ and 3 of the submaximal pfaffians of φ are exactly p_1, p_2, p_3 .*

Proof. Take any alternating map $\psi : F^\vee \rightarrow F$ of rank odd n such that $\text{Pf}_{n-1}(\psi) = I_G = (g_1, \dots, g_n)$, then apply the previous observation to get a new alternating map $\varphi : H^\vee \rightarrow H$ of odd rank $m = n + 6$ such that $\text{Pf}_{m-1}(\varphi) = I_G = (g_1, \dots, g_n, p_1, p_2, p_3, 0, 0, 0)$. $\square$

We need also the following lemma.

Lemma 1.4. *Let M be an alternating matrix of odd size m , with entries in a unitary commutative ring R . Let I be the ideal generated by the submaximal pfaffians of M and let $(q_1, \dots, q_m)$ be a set of generators of I . Then there exists an alternating matrix whose submaximal pfaffians are $uq_1, \dots, uq_m$, where u is a unit of R .*

Proof. Let $\mathbf{p} = (p_1 \dots p_m)$ be the vector of the submaximal pfaffians of M and let $\mathbf{q} = (q_1 \dots q_m)$. Then $\mathbf{p} = \mathbf{q}A$, where A is an invertible square matrix of size m .

Let us consider the alternating matrix $AM^t A$. Now, if N is an alternating matrix we write $\text{pf } N$ for the vector of the submaximal pfaffians of N and if A is a square matrix we write $\bar{A}$ for the adjoint matrix of A . A straightforward computation shows that

$$\text{pf}(AM^t A) = (\text{pf } M)\bar{A} = \mathbf{p}\bar{A} = (\det A)\mathbf{q}.$$

Since A is invertible, $\det A$ is a unit, so we are done. $\square$

Now recall that if $A_G = R/I_G$ is a Gorenstein graded algebra of depth 3 ($R = k[x_0, \dots, x_r]$), its graded minimal free resolution is of the type

$$0 \rightarrow R(-\vartheta_G) \rightarrow \bigoplus_{h \in \vartheta_G - H} R(-h) \rightarrow \bigoplus_{h \in H} R(-h) \rightarrow R \rightarrow A_G \rightarrow 0$$

where H is the multiset of the degrees of the minimal generators of I_G and $\vartheta_G = 2\|H\|/(|H| - 1)$ (remind that $|H|$ must be odd). Hence the multiset H determines all the graded Betti numbers of A_G . Moreover, the integers in $H = \{\{h_1 \leq h_2 \leq \dots \leq h_{2m+1}\}\}$ must satisfy the following Gaeta-Diesel conditions (see [Ga] and [Di]): ϑ_G is an integer and $\vartheta_G > h_{i+1} + h_{2m+2-i}$ for $i = 1, \dots, m$.

In general when T is a graded R -module, (R any polynomial ring), and

$$F_\bullet: 0 \rightarrow \bigoplus_{h \in M_p} R(-h) \rightarrow \dots \rightarrow \bigoplus_{h \in M_{i+1}} R(-h) \rightarrow \bigoplus_{h \in M_i} R(-h) \rightarrow \dots \\ \dots \rightarrow \bigoplus_{h \in M_1} R(-h)$$

is a graded free resolution of T with each M_j a multiset of positive integers, we say $(M_1, \dots, M_p)$ the *multiset sequence* associated to the resolution $F_\bullet$.

If $s \in \text{Supp } M_i \cap \text{Supp } M_{i+1}$, we say that s is a *repetition* in the resolution. Moreover, if $N \subseteq M_i \cap M_{i+1}$ we say that N is *cancellable* for T if there is a graded free resolution of T whose associated multiset sequence is $(M_1, \dots, M_i \setminus N, M_{i+1} \setminus N, \dots, M_p)$.

Next proposition collects some numerical facts about the graded free resolutions of 3-codimensional standard graded Gorenstein algebras.

Proposition 1.5. *Let*

$$0 \rightarrow \bigoplus_{h \in C} R(-h) \rightarrow \bigoplus_{h \in B} R(-h) \rightarrow \bigoplus_{h \in A} R(-h) \rightarrow R \rightarrow R/I_G \rightarrow 0$$

be a graded free resolution (not necessarily minimal) of a depth 3 Gorenstein graded algebra.

- i. $|B \cap C| = |C| - 1$.
- ii. *Let ϑ be the unique element in $\text{Supp}(C \setminus B)$. If $s \in \text{Supp}(A \cap B)$ is such that $\mu_{A \cap B}(s) > \mu_{A \cap B}(\vartheta - s)$ then the multiset $\{s^{(\mu_{A \cap B}(s) - \mu_{A \cap B}(\vartheta - s))}\}$ is cancellable for R/I_G .*

Proof. i. It is enough to remind that in a minimal free resolution of a Gorenstein algebra the module of last syzygies has rank one.

ii. It is enough to remind that in a minimal free resolution of a depth 3 Gorenstein algebra, by selfduality, $\mu_{A \cap B}(s) = \mu_{A \cap B}(\vartheta - s)$. $\square$

Since we need complete intersection ideals contained in Gorenstein ideals, we state some results from the paper [RZ2].

Let

$$\beta = (\{\{d_1, \dots, d_{2n+1}\}\}, \{\{\vartheta - d_{2n+1}, \dots, \vartheta - d_1\}\}, \{\vartheta\}),$$

$n\vartheta = \sum d_i$, $d_1 \leq \dots \leq d_{2n+1}$, be a Betti sequence admissible for an depth 3 standard graded Gorenstein algebra. Let

$$\text{CI}_\beta^g = \{\mathbf{a} \in \mathbb{N}_{\leq}^3 \mid \exists \text{ an ideal } I \subset R \text{ containing a regular sequence of type } \mathbf{a}\}$$

with $\beta_{R/I} = \beta$, R/I Gorenstein ring},

where

$$\mathbb{N}_{\leq}^3 = \{(a_1, a_2, a_3) \in \mathbb{N}^3 \mid a_1 \leq a_2 \leq a_3\}.$$

In [RZ2] Theorem 3.6 it was shown that the poset CI_{β}^g has only one minimal element and it was computed. We report here that statement.

Theorem 1.6. *Let $\beta = (\{\{d_1, \dots, d_{2n+1}\}\}, \{\{\vartheta - d_{2n+1}, \dots, \vartheta - d_1\}\}, \{\vartheta\})$, $n\vartheta = \sum d_i$, $d_1 \leq \dots \leq d_{2n+1}$, be a Betti sequence admissible for an Artinian Gorenstein quotient of $k[x_1, x_2, x_3]$ and define the sets*

$$B = \{3 \leq i \leq n+1 \mid \vartheta \leq d_i + d_{2n+4-i}\}$$

and

$$C = \{4 \leq i \leq n+2 \mid \vartheta \leq d_i + d_{2n+5-i}\}.$$

Then CI_{β}^g has a unique minimal element which we will call $\text{mci } \beta$. Precisely,

- i) if $B \neq \emptyset$, then $\text{mci } \beta = (d_1, d_{\max B}, d_{2n+4-\min B})$;
- ii) if $B = \emptyset$ and $C \neq \emptyset$, then $\text{mci } \beta = (d_1, d_2, d_{\max C})$;
- iii) if $B = \emptyset$ and $C = \emptyset$, then $\text{mci } \beta = (d_1, d_2, d_3)$.

In particular $\text{CI}_{\beta}^g = \{\mathbf{b} \in \mathbb{N}_{\leq}^3 \mid \mathbf{b} \geq \text{mci } \beta\}$.

In the sequel we will need also the following proposition which is a reformulation of Lemma 3.7 of [RZ2].

Proposition 1.7. *Let H be an admissible Hilbert function for an Artinian Gorenstein quotient of $k[x_1, x_2, x_3]$ and let $\beta = (D, \vartheta - D, \{\vartheta\})$ be a Gorenstein Betti sequence compatible with H , with $D = \{\{d_1, \dots, d_{2n+1}\}\}$, $d_1 \leq \dots \leq d_{2n+1}$. Let C be as in Theorem 1.6 and*

$$\overline{B} := \{3 \leq i \leq 2n+1 \mid \vartheta \leq d_i + d_{2n+4-i}\}.$$

We have

- a) $i \in \overline{B} \Rightarrow \mu_D(d_i) = -\Delta^2 H(d_i)$.
- b) $i \in C, i \notin \overline{B}, i-1 \notin \overline{B} \Rightarrow \mu_D(d_i) = -\Delta^2 H(d_i) - 1$.
- c) $i, j \in \overline{B}, d_i = d_j \Rightarrow i = j$.
- d) If $\mu_D(d_i) = -\Delta^2 H(d_i)$ and $k = \min\{j \mid d_j = d_i\}$ then $k \in \overline{B}$.
- e) If $\mu_D(d_i) = -\Delta^2 H(d_i) - 1$ and $k = \min\{j \mid d_j = d_i\} \leq n+2$ then $k \in C$.

Proof. We set $\nu_h := \mu_D(h)$ and $\sigma_h := \mu_{\vartheta-D}(h)$.

- a) Let $i \in \overline{B}$; then $d_i \geq \vartheta - d_{2n+4-i} > d_{i-1}$; furthermore $\vartheta - d_{2n+3-i} > d_i$, so

$$-\Delta^2 H(d_i) = \sum_{h=1}^{d_i} \nu_h - \sum_{h=1}^{d_i} \sigma_h - 1 = (\nu_{d_i} + i - 1) - (i - 2) - 1 = \nu_{d_i}.$$

- b) Let $i \in C \setminus \overline{B}$ such that $i-1 \notin \overline{B}$; then $d_i \geq \vartheta - d_{2n+5-i} > d_{i-1}$; furthermore, since $i \notin \overline{B}$, $\vartheta - d_{2n+4-i} > d_i$, so

$$-\Delta^2 H(d_i) = \sum_{h=1}^{d_i} \nu_h - \sum_{h=1}^{d_i} \sigma_h - 1 = (\nu_{d_i} + i - 1) - (i - 3) - 1 = \nu_{d_i} + 1.$$

c) If $i < j$ then $d_i = d_{i+1}$ so we should have

$$\vartheta \leq d_i + d_{2n+4-i} = d_{i+1} + d_{2n+4-i} < \vartheta.$$

d) We have $d_i > d_1$, therefore $k > 1$ and $d_k > d_{k-1}$; consequently

$$\sum_{h=1}^{d_i-1} \nu_h = \sum_{h=1}^{d_k-1} \nu_h = k-1;$$

moreover by hypotheses

$$-\Delta^2 H(d_i) = \sum_{h=1}^{d_i} \nu_h - \sum_{h=1}^{d_i} \sigma_h - 1 = \nu_{d_i} \Rightarrow \sum_{h=1}^{d_k-1} \nu_h - \sum_{h=1}^{d_k} \sigma_h = 1$$

i.e.

$$\sum_{h=1}^{d_k} \sigma_h = k-2 \Rightarrow \vartheta - d_{2n+4-k} \leq d_k \Rightarrow k \in \overline{B}.$$

e) Again $d_i > d_1$, therefore $k > 1$ and $d_k > d_{k-1}$; consequently

$$\sum_{h=1}^{d_i-1} \nu_h = \sum_{h=1}^{d_k-1} \nu_h = k-1;$$

moreover by hypotheses

$$-\Delta^2 H(d_i) = \sum_{h=1}^{d_i} \nu_h - \sum_{h=1}^{d_i} \sigma_h - 1 = \nu_{d_i} + 1 \Rightarrow \sum_{h=1}^{d_k-1} \nu_h - \sum_{h=1}^{d_k} \sigma_h = 2$$

i.e.

$$\sum_{h=1}^{d_k} \sigma_h = k-3 \Rightarrow \vartheta - d_{2n+5-k} \leq d_k,$$

hence $k \in C$.

□

Remark 1.8. We remind that by [RZ1], Proposition 3.7, for every $d_i > d_1$, $-\Delta^2 H(d_i)$ is equal to the largest number of minimal generators of degree d_i (which we will denote by $\text{Mng}_H(d_i)$) compatible with an Artinian Gorenstein graded algebra of codimension 3 having Hilbert function equal to H .

Remark 1.9. Note that if $B = \emptyset$ then $\overline{B} \subseteq \{n+2\}$. Indeed let $i \in \overline{B}$; since $B = \emptyset$ we have that $i \geq n+2$; moreover $\vartheta \leq d_i + d_{2n+4-i}$, so $2n+4-i \geq n+2$, hence $i \leq n+2$ i.e. $i = n+2$.

2. STRUCTURE THEOREM FOR ALMOST COMPLETE INTERSECTIONS

In the sequel, if $f : M \rightarrow N$ and $g : M \rightarrow P$ are maps of modules we will write $(f, g) : M \rightarrow N \oplus P$ for the map defined by $(f, g)(m) = (f(m), g(m))$. Moreover, if $f : M \rightarrow P$ and $g : N \rightarrow P$ we will write $f|g : M \oplus N \rightarrow P$ for the map defined by $f|g(m, n) = f(m) + g(n)$.

Let R be a Noetherian local commutative ring, H_0 a free R -module of odd rank $m_0 \geq 5$ and $\varphi_0 : H_0^\vee \rightarrow H_0$ an alternating map. Let $J := \text{Pf}_{m_0-1}(\varphi_0)$ be the ideal generated by the pfaffians of φ_0 of size $m_0 - 1$ (note that despite the fact that the pfaffians depend on the choice of the

bases in H_0 , J depends only on the map φ_0). Moreover, we denote by $\text{pf } \varphi_0 : H_0 \rightarrow R$ the map defined by the submaximal pfaffians of φ_0 . It is known by Buchsbaum-Eisenbud Theorem (see [BE]) that $\text{depth } J \leq 3$ and we suppose here that $\text{depth } J = 3$. Take a regular sequence (p_1, p_2, p_3) in J . By Lemma 1.3 there exists an alternating map $\varphi : H^\vee \rightarrow H$ of odd rank m such that $\text{Pf}_{m-1}(\varphi) = J$ and 3 of the submaximal pfaffians of φ are exactly p_1, p_2, p_3 . So $J = (p_1, p_2, p_3, \dots, p_m)$, where the p_i 's, for $1 \leq i \leq m$, are the submaximal pfaffians of φ .

Let $I = (p_1, p_2, p_3)$ and $\psi : G^\vee \rightarrow G$ an alternating map whose pfaffians are exactly p_1, p_2, p_3 . Then $H = G \oplus F$, where F is a free R -module of rank $m-3$, and $\text{pf } \varphi = \text{pf } \psi | \sigma$, where $\sigma : F \rightarrow R$. Therefore we have the following decomposition: $\varphi = (\alpha | -\lambda^\vee, \lambda | \beta)$, where $\alpha : G^\vee \rightarrow G$, $\beta : F^\vee \rightarrow F$ and $\lambda : G^\vee \rightarrow F$. Moreover we denote by $\bar{\beta} : F \rightarrow F^\vee$, the alternating map such that $\beta \bar{\beta} = p \text{ id}$ and $\bar{\beta} \beta = p \text{ id}$, where p is the pfaffian of the map β . One can see that (with respect to suitable bases) the matrix associated to $\bar{\beta}$ is the pfaffian adjoint of the matrix of β .

[The pfaffian of an empty matrix will be 1; note that $\text{pf } \psi^\vee = -(\text{pf } \psi)$]

Proposition 2.1. *With the above notation*

$$0 \longrightarrow F^\vee \xrightarrow{(\lambda^\vee, -\beta)} G \oplus F \xrightarrow{(p | \lambda^\vee \bar{\beta}, -\text{pf } \psi | -\sigma)} G \oplus R \xrightarrow{\text{pf } \psi | p} R$$

is a free resolution of an almost complete intersection algebra R/Q .

Proof. We start by proving that the canonical surjection $R/I \rightarrow R/J$ can be lifted to a map between their resolutions in the following way:

$$\begin{array}{ccccccc} 0 & \longrightarrow & R & \xrightarrow{(\text{pf } \psi)^\vee} & G^\vee & \xrightarrow{\psi} & G & \xrightarrow{\text{pf } \psi} & R \\ & & \downarrow p & & \downarrow (p, -\bar{\beta}\lambda) & & \downarrow (\text{id}, 0) & & \downarrow \text{id} \\ 0 & \longrightarrow & R & \xrightarrow{((\text{pf } \psi)^\vee, \sigma^\vee)} & G^\vee \oplus F^\vee & \xrightarrow{(\alpha | -\lambda^\vee, \lambda | \beta)} & G \oplus F & \xrightarrow{\text{pf } \psi | \sigma} & R \end{array}$$

Indeed, since

$$\begin{aligned} 0 &= (\lambda | \beta)((\text{pf } \psi)^\vee, \sigma^\vee) = \lambda(\text{pf } \psi)^\vee + \beta\sigma^\vee \Rightarrow \\ &\Rightarrow \bar{\beta}\lambda(\text{pf } \psi)^\vee + \bar{\beta}\beta\sigma^\vee = 0 \Rightarrow \bar{\beta}\lambda(\text{pf } \psi)^\vee + p\sigma^\vee = 0 \end{aligned}$$

we have

$$(p, -\bar{\beta}\lambda)(\text{pf } \psi)^\vee = (p(\text{pf } \psi)^\vee, -\bar{\beta}\lambda(\text{pf } \psi)^\vee) = (p(\text{pf } \psi)^\vee, p\sigma^\vee) = p((\text{pf } \psi)^\vee, \sigma^\vee).$$

Now Remark 1.2 will imply, in our notation, that $\alpha p + \lambda^\vee \bar{\beta}\lambda = \psi$. Therefore we have

$$\begin{aligned} (\alpha | -\lambda^\vee, \lambda | \beta)(p, -\bar{\beta}\lambda) &= \\ &= (\alpha p + \lambda^\vee \bar{\beta}\lambda, \lambda p - \beta \bar{\beta}\lambda) = (\psi, \lambda p - p\lambda) = (\psi, 0) = (\text{id}, 0)\psi. \end{aligned}$$

Then we set $\mathbb{F}_\bullet^I$ and $\mathbb{F}_\bullet^J$ the above resolutions of R/I and R/J and $\tau : \mathbb{F}_\bullet^I \rightarrow \mathbb{F}_\bullet^J$ the above complex map. Thus, if we set $Q := I : J$, we see that a free resolution of R/Q is given by the mapping cone of the map

$\tau^\vee : (\mathbb{F}_\bullet^J)^\vee \rightarrow (\mathbb{F}_\bullet^I)^\vee$. If we make cancellations where we have identities map we get the required resolution of the almost complete intersection R/Q . $\square$

As a simple consequence of the previous result we have

Corollary 2.2. *Let R be a Noetherian local ring and let M be an alternating matrix of odd rank m , whose entries are in R . Let $p_1, \dots, p_m$ be the submaximal pfaffians of M and p_{abc} the pfaffian of order $m-3$ obtained by M by deleting the rows and columns a, b, c . If (p_a, p_b, p_c) is a regular sequence, then*

$$(p_a, p_b, p_c) : (p_1, \dots, p_m) = (p_a, p_b, p_c, p_{abc}).$$

The Proposition 2.1 gives a way to construct almost complete intersection algebras but the very interesting thing is that every almost complete intersection can be constructed in this way.

Theorem 2.3. *Let R be a Noetherian local ring and $I_Q \subset R$ be a perfect ideal of depth 3 of an almost complete intersection. Then there exists an alternating map $\psi : H^\vee \rightarrow H$, where H is a free R -module of odd rank with $\text{impf } \psi$ of depth 3, such that, with the above notation,*

$$0 \longrightarrow F^\vee \xrightarrow{(\lambda^\vee, -\beta)} G \oplus F \xrightarrow{(p|\lambda^\vee \bar{\beta}, -\text{pf } \psi|-\sigma)} G \oplus R \xrightarrow{\text{pf } \psi|p} R$$

is a free resolution of R/I_Q .

Proof. Since I_Q is an almost complete intersection ideal we have that $I_Q = (p_0, p_1, p_2, p_3)$ and since $\text{depth } I_Q = 3$ we can suppose that $I_Z := (p_1, p_2, p_3)$ is generated by a regular sequence. Let $I_G := I_Z : I_Q$. Then I_G is a Gorenstein ideal of depth 3. By Lemma 1.3, there exists an alternating map $\varphi : H^\vee \rightarrow H$, such that the submaximal pfaffians of φ are exactly p_1, p_2, p_3 and the other generators of I_G . By Proposition 2.1 the above resolution gives a resolution of R/I_Q . $\square$

Now we would like to give a graded version of the previous result.

Let $\mathbb{P}_k^r$ be the projective space with $r \geq 3$ and $R = k[x_0, x_1, \dots, x_r]$ the standard graded coordinate ring of $\mathbb{P}_k^r$.

Theorem 2.4. *Let $Q \subset \mathbb{P}_k^r$ be an almost complete intersection scheme of codimension 3 and $I_Q \subset R$ be its defining ideal. Then R/I_Q admits a graded free resolution of the following type*

$$0 \rightarrow K^\vee(-d) \rightarrow G(-d_0) \oplus K \rightarrow G \oplus R(-d_0) \rightarrow R$$

where d_0 is a positive integer, $G = \bigoplus_{i=1}^3 R(-d_i)$, $d = d_0 + d_1 + d_2 + d_3$, $K = \bigoplus_{i=4}^m R(-e_i)$, $m \geq 5$ an odd integer and $d_1, d_2, d_3, e_4 - d_0, \dots, e_m - d_0$ are the degrees of all submaximal pfaffians of a suitable alternating matrix of size m .

Proof. Let $Z \subset \mathbb{P}_k^r$ be a complete intersection generated by 3 minimal generators of I_Q , of degrees d_1, d_2, d_3 . Let $I_\Gamma := I_Z : I_Q$. I_Γ is the saturated homogeneous ideal of an aG scheme Γ directly linked to Q in Z . As in

Proposition 2.1 we have the following diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & R(-\vartheta_Z) & \longrightarrow & G^\vee(-\vartheta_Z) & \longrightarrow & G \longrightarrow R \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & R(-\vartheta_G) & \longrightarrow & G^\vee(-\vartheta_G) \oplus F^\vee(-\vartheta_G) & \xrightarrow{\varphi} & G \oplus F \longrightarrow R
\end{array}$$

where the first row is a minimal graded resolution of R/I_Z and the second row is the graded resolution of R/I_Γ , obtained by the alternating map φ as in Lemma 1.3. So we have that $G = \oplus_{i=1}^3 R(-d_i)$ and $F = \oplus_{i=4}^m R(-d_i)$. Dualizing and shifting by $-\vartheta_Z$ this diagram we get

$$\begin{array}{ccccccc}
0 \longrightarrow & R(-\vartheta_Z) & \longrightarrow & G^\vee(-\vartheta_Z) \oplus F^\vee(-\vartheta_Z) & \longrightarrow & G(-d_0) \oplus F(-d_0) & \longrightarrow R(-d_0) \\
& \downarrow & & \downarrow & & \downarrow & \downarrow \\
0 \longrightarrow & R(-\vartheta_Z) & \longrightarrow & G^\vee(-\vartheta_Z) & \longrightarrow & G & \longrightarrow R
\end{array}$$

where $d_0 := \vartheta_Z - \vartheta_G$. Taking the mapping cone and after the trivial cancellations we obtain the following resolution of R/I_Q

$$0 \rightarrow F^\vee(-\vartheta_Z) \rightarrow G(-d_0) \oplus F(-d_0) \rightarrow G \oplus R(-d_0) \rightarrow R.$$

Now if we set $K := F(-d_0)$ and $d := 2\vartheta_Z - \vartheta_G = d_0 + d_1 + d_2 + d_3$ we get

$$0 \rightarrow K^\vee(-d) \rightarrow G(-d_0) \oplus K \rightarrow G \oplus R(-d_0) \rightarrow R,$$

which is the required resolution. $\square$

Corollary 2.5. *Let $Q \subset \mathbb{P}_k^r$ be an almost complete intersection scheme of codimension 3 and $I_Q \subset R$ be its defining ideal, then R/I_Q admits a graded free resolution of the type*

$$0 \rightarrow K^\vee(-d) \rightarrow G(-d_0) \oplus K \rightarrow G \oplus R(-d_0) \rightarrow R$$

in which $d_0 = \min\{\deg p \mid p \in I_Q\}$.

Proof. Let $d_0 \leq d_1 \leq d_2 \leq d_3$ the degrees of a minimal set of generators of I_Q . Since we can find a regular sequence of minimal generators in I_Q of type d_1, d_2, d_3 , the conclusion follows by repeating the argument of Theorem 2.4 using such a regular sequence. $\square$

Example 2.6. Let $Q \subset \mathbb{P}_k^3$ be the 0-dimensional almost complete intersection linked to 5 general points (Γ) in a complete intersection Z of type $(2, 2, 8)$. Then we consider the following graded resolutions

$$0 \rightarrow R(-12) \rightarrow R(-4) \oplus R(-10)^2 \rightarrow R(-2)^2 \oplus R(-8) \rightarrow R \rightarrow R/I_Z \rightarrow 0$$

and

$$\begin{aligned}
0 \rightarrow R(-5) \rightarrow R(-8) \oplus R(-3)^5 \oplus R(3) \rightarrow R(-8) \oplus R(-2)^5 \oplus R(3) \rightarrow \\
\rightarrow R \rightarrow R/I_\Gamma \rightarrow 0,
\end{aligned}$$

where the last one is the pfaffian resolution of R/I_Γ got by the minimal one by adding the term $R(-8) \oplus R(3)$ to the second and to the third module of the complex. Such a resolution can be built as illustrated in Lemma 1.3 and the pfaffians of its alternating central map are the five minimal generators

of I_Γ of degree 2, the form of I_Γ of degree 8 used to perform the linkage and the null form. So we have $G = R(-2)^2 \oplus R(-8)$, $K = R(-9)^3 \oplus R(-4)$, $d_0 = 7$ and $d = 19$. Consequently we get the following resolution of R/I_Q

$$\begin{aligned} 0 \rightarrow R(-15) \oplus R(-10)^3 \rightarrow [R(-9)^2 \oplus R(-15)] \oplus [R(-9)^3 \oplus R(-4)] \rightarrow \\ \rightarrow [R(-8) \oplus R(-2)^2] \oplus R(-7) \rightarrow R \rightarrow R/I_Q \rightarrow 0. \end{aligned}$$

Note that a minimal graded resolution of R/I_Q can be obtained by deleting the term $R(-15)$.

3. THE GRADED BETTI NUMBERS OF ALMOST COMPLETE INTERSECTIONS

In this section we would like to characterize all graded Betti sequences admissible for an almost complete intersection scheme of codimension 3 of $\mathbb{P}^r$.

Since we are interested to the graded Betti numbers for such schemes we can restrict ourselves to the Artinian reduction of X . So, from now on, we let $R = k[x_1, x_2, x_3]$ and $A_Q = R/I_Q$ an Artinian almost complete intersection graded algebra. A graded minimal free resolution of A_Q is of the following form

$$0 \rightarrow \bigoplus_{h \in F} R(-h) \rightarrow \bigoplus_{h \in E} R(-h) \rightarrow \bigoplus_{h \in D} R(-h) \rightarrow R \rightarrow A_Q \rightarrow 0$$

where D, E, F are multisets of positive integers with $|D| = 4$, $|E| = |F| + 3$.

For this aim we will consider the following multisets of positive integers $D = \{\{d_0 \leq d_1 \leq d_2 \leq d_3\}\}$, $E = \{\{e_i\}_{i=1}^{p+3}\}$ and $F = \{\{f_i\}_{i=1}^p\}$ where $p \geq 2$. In the sequel we will denote $D^* = \{\{d_1 \leq d_2 \leq d_3\}\}$.

Lemma 3.1. *Let (D, E, F) be a Betti sequence admissible for an Artinian almost complete intersection graded algebra of codimension 3. Then*

- 1) $d - F \subset E$, where $d = \|D\|$;
- 2) if we set $\widehat{E} := E \setminus (d - F)$, $S := D^* \cap (\vartheta_Z - \widehat{E})$, $\overline{D} := D^* \setminus S$ then $\widehat{E} = (d_0 + \overline{D}) \sqcup (\vartheta_Z - S)$.

Proof. 1) Let $A = R/I_Q$ be an Artinian almost complete intersection graded algebra such that $\beta_A = (D, E, F)$. I_Q contains a length 3 regular sequence of type (d_1, d_2, d_3) which is part of a minimal set of generators for I_Q ; let I_Z be the ideal generated by such a regular sequence. Then $I_G := I_Z : I_Q$ is a Gorenstein ideal of depth 3. By standard mapping cone procedure we obtain the following graded free resolution of R/I_G

$$0 \rightarrow R(-\vartheta_G) \rightarrow \bigoplus_{h \in \vartheta_Z - E} R(-h) \rightarrow \bigoplus_{h \in \vartheta_Z - F} R(-h) \oplus \bigoplus_{h \in D^*} R(-h) \rightarrow R$$

where $D^* = \{\{d_1, d_2, d_3\}\}$, $\vartheta_Z = \|D^*\|$, $\vartheta_G = \vartheta_Z - d_0$. Since, by mapping cone, no summand of $\bigoplus_{h \in \vartheta_Z - F} R(-h)$ is cancellable in such a resolution, by Gorenstein duality, $\vartheta_G - (\vartheta_Z - F) \subset \vartheta_Z - E$ hence $\vartheta_Z - (\vartheta_G - (\vartheta_Z - F)) \subset \vartheta_Z - (\vartheta_Z - E)$, i.e. $d - F \subset E$.

2) By our setting we have $D^* = S \sqcup \overline{D}$ and $\vartheta_Z - \widehat{E} = S \sqcup [(\vartheta_Z - \widehat{E}) \setminus S]$; therefore in the previous resolution we have that

$$\begin{aligned} \bigoplus_{h \in \vartheta_Z - E} R(-h) &= \bigoplus_{h \in S} R(-h) \oplus \bigoplus_{h \in (\vartheta_Z - \widehat{E}) \setminus S} R(-h) \oplus \bigoplus_{h \in -d_0 + F} R(-h) \\ \bigoplus_{h \in D^*} R(-h) &= \bigoplus_{h \in S} R(-h) \oplus \bigoplus_{h \in \overline{D}} R(-h) \end{aligned}$$

where the only summands eventually cancellable are in $\bigoplus_{h \in S} R(-h)$, therefore, by Gorenstein duality, $\vartheta_G - \overline{D} = (\vartheta_Z - \widehat{E}) \setminus S$, hence $\vartheta_Z - (\vartheta_G - \overline{D}) = \vartheta_Z - [(\vartheta_Z - \widehat{E}) \setminus S]$, i.e. $d_0 + \overline{D} = \widehat{E} \setminus (\vartheta_Z - S)$ and we are done. $\square$

Because of the previous lemma it is convenient to give the following definition.

Definition 3.2. A sequence (D, E, F) of multisets of positive integers is said of *aci*-type if it satisfies the following conditions

- 1) $|D| = 4$, $|E| = |F| + 3$, $|F| \geq 2$;
- 2) $d - F \subset E$, where $d := \|D\|$;
- 3) if we set $\widehat{E} := E \setminus (d - F)$, $d_0 := \min D$, $D^* := D \setminus \{d_0\}$, $\vartheta_Z = \|D^*\|$, $S := D^* \cap (\vartheta_Z - \widehat{E})$, $\overline{D} := D^* \setminus S$ then $\widehat{E} = (d_0 + \overline{D}) \sqcup (\vartheta_Z - S)$.

Remark 3.3. Let $A = R/I_Q$ be an Artinian almost complete intersection graded algebra such that $\beta_A = (D, E, F)$; let I_Z be an ideal generated by a regular sequence of type (d_1, d_2, d_3) which is part of a minimal set of generators for I_Q and $I_G := I_Z : I_Q$ the linked Gorenstein ideal of depth 3. Then the minimal graded free resolution of $A_G = R/I_G$ has the following form

$$\begin{array}{ccccc} \bigoplus_{h \in \vartheta_Z - F} R(-h) & \oplus & \bigoplus_{h \in \overline{D}} R(-h) & \oplus & \bigoplus_{h \in \overline{S}} R(-h) \\ & & \uparrow & & \\ \bigoplus_{h \in -d_0 + F} R(-h) & \oplus & \bigoplus_{h \in \vartheta_G - \overline{D}} R(-h) & \oplus & \bigoplus_{h \in \overline{S}} R(-h) \\ & & \uparrow & & \\ & & R(-\vartheta_G) & & \\ & & \uparrow & & \\ & & 0 & & \end{array}$$

where $\overline{S} \subseteq S$. Now observe that $\vartheta_G - [(\vartheta_Z - F) \sqcup \overline{D}] = (-d_0 + F) \sqcup (\vartheta_G - \overline{D})$, so we can apply Proposition 1.1 and deduce that for $\overline{S}$ there are 4 possibilities depending on its cardinality.

- 0) $\overline{S} = \emptyset$;
- 1) $|\overline{S}| = 1$ in such a case, by Proposition 3.7 and Remark 3.8 in [RZ1], $\overline{S} = \{\vartheta_G/2\}$;
- 2) $|\overline{S}| = 2$ in such a case, by Proposition 3.7 and Remark 3.8 [RZ1], $\overline{S} = \{\{\alpha, \vartheta_G - \alpha\}\}$ for some α ;
- 3) $|\overline{S}| = 3$ in such a case, by Proposition 3.7 and Remark 3.8 [RZ1], $\overline{S} = \{\{\vartheta_G/2, \alpha, \vartheta_G - \alpha\}\}$ for some α .

In particular, if $\vartheta_G/2 \notin \text{Supp } S$ then $|\overline{D}| + |F|$ is odd and consequently $|S| + |F|$ is even. It is easy to produce examples in which $\vartheta_G/2 \in \text{Supp } S$ and $|S| + |F|$ is even and examples in which $\vartheta_G/2 \in \text{Supp } S$ and $|S| + |F|$ is odd.

Now we prove two technical lemmas which will be crucial for the characterization of the Betti sequences of almost complete intersections.

Lemma 3.4. *Let A_G be an Artinian Gorenstein graded algebra of codimension 3 whose Betti sequence is β_G . Let $(d_1, d_2, d_3) \geq (e_1, e_2, e_3) := \text{mci } \beta_G$, with $d_1 \leq d_2 \leq d_3$ and $e_1 \leq e_2 \leq e_3$.*

- 1) *If $d_i = e_i$ for some i then for every regular sequence (g_1, g_2, g_3) in I_G , with $\deg g_j = d_j$ for $1 \leq j \leq 3$, g_i is a minimal generator for I_G .*
- 2) *If $d_i > e_i$ for some i and I_G has a minimal generator of degree d_i then there are in I_G regular sequences (g_1, g_2, g_3) , with $\deg g_j = d_j$ for $1 \leq j \leq 3$, such that g_i is a minimal generator for I_G and regular sequences (h_1, h_2, h_3) , with $\deg h_j = d_j$ for $1 \leq j \leq 3$, such that h_i is not a minimal generator for I_G .*

Proof. 1) Let us suppose that $d_i = e_i$ for some i and let us consider a regular sequence (g_1, g_2, g_3) in I_G , with $\deg g_j = d_j$ for $1 \leq j \leq 3$. It is enough to remind that if $i = 1$ then $d_1 = \min\{\deg f \mid f \in I_G\}$; if $i = 2$ then $\text{depth}(I_G)_{\leq d_2-1} = 1$; if $i = 3$ then $\text{depth}(I_G)_{\leq d_3-1} = 2$.

2) Let us suppose that $d_i > e_i$ for some i and that I_G has a minimal generator of degree d_i . Let (f_1, f_2, f_3) be a regular sequence in I_G such that $\deg f_j = e_j$ for $1 \leq j \leq 3$. If we choose $h_j := f_j a_j$, where a_j is a generic form of degree $d_j - e_j$, for $1 \leq j \leq 3$, we get a regular sequence in which h_i is not a minimal generator for I_G . Now we set $g_j := f_j a_j$, where a_j is a generic form of degree $d_j - e_j$ for $j \neq i$ and we take as g_i a generic form in $(I_G)_{d_i}$. Of course, (g_1, g_2, g_3) is a regular sequence and, since by hypothesis I_G has minimal generators in degree d_i , g_i will be a minimal generator for I_G . $\square$

Lemma 3.5. *Let $A_\Gamma = R/I_\Gamma$ be an Artinian Gorenstein graded algebra of codimension 3, whose last syzygy degree is ϑ_Γ . Let (f_1, f_2, f_3) be a regular sequence in I_Γ , with $\deg f_i = d_i$ and $d_1 \leq d_2 \leq d_3$. Let us suppose that $d_i + d_j = \vartheta_\Gamma$ and f_i, f_j are minimal generators for I_Γ for some $1 \leq i < j \leq 3$. Let A_G be an Artinian Gorenstein graded algebra of codimension 3 whose Betti sequence is obtained from the Betti sequence of A_Γ by deleting the degrees d_i and d_j between generators and first syzygies. If $(e_1, e_2, e_3) := \text{mci } \beta_G$, then $d_i > e_i$ and $d_j > e_j$.*

Proof. Along this proof we let $\gamma_1 \leq \dots \leq \gamma_{2m+1}$ be the degrees of a minimal set of generators for I_Γ , $g_1 \leq \dots \leq g_{2m-1}$ be the degrees of a minimal set of generators for I_G and $(\epsilon_1, \epsilon_2, \epsilon_3) := \text{mci } \beta_\Gamma$.

We start with the more delicate case, i.e. for $i = 2$ and $j = 3$. By [RZ3] Theorem 3.9, we have $\text{mci } \beta_G \leq \text{mci } \beta_\Gamma \leq (d_1, d_2, d_3)$, so if $\epsilon_2 < d_2$ the conclusion follows for d_2 . Therefore we can suppose that $\epsilon_2 = d_2$. If A is an Artinian Gorenstein graded algebra of codimension 3, whose Hilbert function is H , we write $\text{Mng}_H(n)$ as the largest number of minimal generators of degree n compatible with an Artinian Gorenstein graded algebra of codimension 3 having Hilbert function equal to H (see Remark 1.8).

Now, if H is the Hilbert function of A_Γ , according to Proposition 1.8 and Theorem 1.6, either $\epsilon_2 = \gamma_2$ or I_Γ has $\text{Mng}_H(\epsilon_2)$ minimal generators of degree ϵ_2 . But $\epsilon_2 = \gamma_2$ implies that $\gamma_2 = \epsilon_2 = d_2 = \vartheta_\Gamma - d_3$ is a degree of a first syzygy of I_Γ which is clearly impossible. So, I_Γ has $\text{Mng}_H(\epsilon_2)$ minimal generators of degree $\epsilon_2 = d_2$. On the other hand the Hilbert function of A_G is still H and I_G has $\text{Mng}_H(d_2) - 1$ minimal generators in degree d_2 . This implies, again by Proposition 1.8 and Theorem 1.6, that $e_2 < \epsilon_2 = d_2$.

Now, if $d_3 > \epsilon_3$ we are done. So we can assume that $d_3 = \epsilon_3$. Observe that first $\epsilon_3 > \gamma_3$: indeed, otherwise, $d_3 = \epsilon_3 = \gamma_3$ and consequently $d_2 = \epsilon_2 = \gamma_2$ (we remind that d_2 and d_3 are degrees of minimal generators for I_Γ) and we should have again a first syzygy of degree γ_2 for I_Γ . So we have two possibilities: either $\epsilon_2 = \gamma_2$ or $\epsilon_2 > \gamma_2$. If $\epsilon_2 = \gamma_2$, since $\epsilon_3 > \gamma_3$, by Theorem 1.6 $B_\Gamma = \emptyset$ and $C_\Gamma \neq \emptyset$. If $\overline{B}_\Gamma = \emptyset$ then I_Γ has $\text{Mng}_H(\epsilon_3) - 1$ minimal generators in degree ϵ_3 . Then $e_2 = \gamma_2$ and I_G has $\text{Mng}_H(\epsilon_3) - 2$ generators in degree ϵ_3 ; therefore $e_3 < \epsilon_3$, i.e. $d_3 > e_3$. If $\overline{B}_\Gamma = \{n+2\}$ (see Remark 1.9) then $\gamma_{n+2} \geq \vartheta_\Gamma - \gamma_{n+2} > \gamma_{n+1}$ hence $e_3 \leq g_{n+1} = \gamma_{n+1} < \gamma_{n+2} = \epsilon_3$ (note that, in this case, $g_i = \gamma_i$, for $i \leq n+1$).

Finally, if $\epsilon_2 > \gamma_2$ then

$$B_\Gamma = \{3 \leq h \leq m+1 \mid \vartheta_\Gamma \leq \gamma_h + \gamma_{2m+4-h}\} \neq \emptyset$$

(as defined in Theorem 1.6) and consequently I_Γ has $\text{Mng}_H(d_3)$ minimal generators in degree d_3 . Let

$$B_G = \{3 \leq h \leq m \mid \vartheta_\Gamma \leq g_h + g_{2m+2-h}\}$$

and

$$C_G = \{4 \leq h \leq m+1 \mid \vartheta_\Gamma \leq g_h + g_{2m+3-h}\}$$

Now, if $B_G \neq \emptyset$, since I_G has $\text{Mng}_H(d_3) - 1$ minimal generators in degree d_3 , we see that $e_3 < \epsilon_3 = d_3$. If $B_G = \emptyset$ and we set $b := \max B_\Gamma$, there exist only two integers, γ_b and γ_{2m+4-b} , in which I_Γ has the maximum number of minimal generators allowed by H . Therefore $\vartheta_\Gamma - \gamma_b = \gamma_{2m+4-b} = \epsilon_3 = d_3$ and this implies also that $d_2 = \gamma_b = \epsilon_2$. Moreover $g_h = \gamma_h$ for $1 \leq h \leq b-1$, $g_h = \gamma_{h+1}$ for $b \leq h \leq 2m+2-b$ and $g_h = \gamma_{h+2}$ for $2m+3-b \leq h \leq 2m-1$. Therefore

$$g_b = \gamma_{b+1} \geq \gamma_b = \vartheta_\Gamma - \gamma_{2m+4-b} \geq \vartheta_\Gamma - \gamma_{2m+5-b} = \vartheta_\Gamma - g_{2m+3-b}$$

i.e. $b \in C_G$. Let $c := \max C_G$. Of course $b \leq c \leq m+1 \leq 2m+2-b$, so $e_3 = g_c = \gamma_{c+1}$. Therefore we have $g_c = \gamma_{c+1} = e_3 \leq \epsilon_3 = d_3 = \gamma_{2m+4-b}$. Suppose that $e_3 = \epsilon_3 = d_3$, i.e. $\gamma_{c+1} = \gamma_{2m+4-b}$. Since $c+1 \leq m+2 < 2m+4-b$, we have $\gamma_h = \gamma_{2m+4-b}$ for $c+1 \leq h < 2m+4-b$. Consequently, $\gamma_h + \gamma_{2m+4-h} \geq \gamma_{2m+4-b} + \gamma_b = \vartheta_\Gamma$, i.e. h and $2m+4-h$ satisfy the inequality to stay in B_Γ ; but $h > b$ and $2m+4-h > b$ therefore $h \notin B_\Gamma$ and $2m+4-h \notin B_\Gamma$ so we get that $h \geq m+2$ and $2m+4-h \geq m+2$ i.e. $h = m+2$. So we are reduced to the case $c+1 = m+2$ and $2m+4-b = m+3$ i.e. $c = b = m+1$ and $\gamma_{m+2} = \gamma_{m+3}$. So we have that

$$\vartheta_\Gamma = \gamma_b + \gamma_{2m+4-b} = \gamma_{m+1} + \gamma_{m+3} = \gamma_{m+1} + \gamma_{m+2} < \vartheta_\Gamma$$

where the last inequality holds for the Gaeta-Diesel conditions. So we get a contradiction and consequently $e_3 < \epsilon_3 = d_3$.

The remaining cases are simpler. So, if $i = 1$ and $j = 2$ since $\vartheta_\Gamma - d_2 = d_1$ is

a first syzygy for I_Γ we have $d_1 > \epsilon_1 \geq e_1$. Moreover, $d_2 = \vartheta_\Gamma - d_1 \geq \vartheta_\Gamma/2 \geq \gamma_{m+1} \geq \epsilon_2$. thus, if $d_2 = \epsilon_2$ we should have $d_2 = \vartheta_\Gamma/2 = d_1 = \gamma_{m+1}$ and, consequently, $\vartheta_\Gamma/2 = \vartheta_\Gamma - \gamma_{m+1} > \gamma_{m+2} \geq \vartheta_\Gamma/2$, a contradiction, therefore $d_2 > \epsilon_2 \geq e_2$.

Finally, if $i = 1$ and $j = 3$, as before we easily get that $d_1 > \epsilon_1 \geq e_1$. Of course, we can suppose $d_3 = \epsilon_3$. We then show that $B_\Gamma \neq \emptyset$: namely, otherwise, $d_3 = \epsilon_3 \leq \gamma_{m+2}$, let us say $d_3 = \gamma_h$ for some $h \leq m+2$. Thus we should have

$$d_1 = \vartheta_\Gamma - d_3 = \vartheta_\Gamma - \gamma_h > \gamma_{2m+3-h} \geq \gamma_{m+1} \geq d_2$$

a contradiction. Therefore, we have $B_\Gamma \neq \emptyset$ and as in the case $i = 2$ and $j = 3$ we are reduced to the case in which B_Γ has only one element b . In this case we have $\epsilon_2 = \gamma_b$ and $\epsilon_3 = d_3 = \gamma_{2m+4-b}$; then $d_1 = \vartheta_\Gamma - d_3 = \vartheta_\Gamma - \gamma_{2m+4-b} > \gamma_{b-1}$ and $\gamma_b \geq \vartheta_\Gamma - d_3 = d_1$ since $b \in B_\Gamma$, i.e. $d_1 = \gamma_b$ and the conclusion $e_3 < \epsilon_3 = d_3$ follows as in the final part of the previous case $i = 2$ and $j = 3$. $\square$

Theorem 3.6. *Let (D, E, F) be a sequence of multisets of positive integers. (D, E, F) is a Betti sequence admissible for an Artinian almost complete intersection algebra of codimension 3 if and only if it satisfies the following conditions:*

- 1) *it is an aci-type sequence;*
- 2) *$\beta_G := (G_0, G_1, G_2)$ is the Betti sequence of a 3-codimensional Gorenstein Artinian graded algebra, where*

$$G_0 := (\vartheta_Z - F) \sqcup \overline{D} \sqcup T$$

$$G_1 := (-d_0 + F) \sqcup (\vartheta_G - \overline{D}) \sqcup T$$

$$G_2 := \{\vartheta_G\}$$

with $\vartheta_G := \vartheta_Z - d_0$ and $T := \{\vartheta_G/2\}$ when $\vartheta_G/2 \in \text{Supp } S$ and $|F| + |\overline{D}|$ is even and $T := \emptyset$ otherwise.

- 3) *if we set $(e_1, e_2, e_3) := \text{mci } \beta_G$, $e_1 \leq e_2 \leq e_3$, $D^* = \{\{d_1, d_2, d_3\}\}$, $d_1 \leq d_2 \leq d_3$ then $d_i \geq e_i$ for $1 \leq i \leq 3$ and for every $s \in \text{Supp}(S \setminus T)$, $d_i > e_i$ for $i := \min\{j \mid d_j = s\} + \mu_{S \setminus T}(s) - 1$.*

Proof. Let us suppose that (D, E, F) is the Betti sequence of an almost complete intersection of codimension 3. By Lemma 3.1 (D, E, F) is an aci-type sequence. Using the same terminology of the Definition 3.2 we denote $D = \{\{d_0, d_1, d_2, d_3\}\}$ with $d_0 \leq d_1 \leq d_2 \leq d_3$. Let $A_Q = R/I_Q$ be an almost complete intersection Artinian graded algebra with Betti sequence (D, E, F) and $A_Z = R/I_Z$ be a complete intersection Artinian graded algebra of type (d_1, d_2, d_3) with I_Z generated by f_1, f_2, f_3 minimal generators of I_Q . Note that such A_Z must exist since $d_1, d_2, d_3 \geq d_0$. Let $I_\Gamma := I_Z : I_Q$. Of course, $A_\Gamma = R/I_\Gamma$ is a Gorenstein Artinian graded algebra. Using Remark 3.3 a

minimal graded free resolution of A_Γ will be of the type

$$\begin{array}{c}
 \bigoplus_{h \in \vartheta_Z - F} R(-h) \oplus \bigoplus_{h \in \overline{D}} R(-h) \oplus \bigoplus_{h \in \overline{S}} R(-h) \\
 \uparrow \\
 \bigoplus_{h \in -d_0 + F} R(-h) \oplus \bigoplus_{h \in \vartheta_G - \overline{D}} R(-h) \oplus \bigoplus_{h \in \overline{S}} R(-h) \\
 \uparrow \\
 R(-\vartheta_\Gamma) \\
 \uparrow \\
 0
 \end{array}$$

Using Remark 3.3 one sees that either $\overline{S} = T$ or $\overline{S} \setminus T = \{\{\alpha, \vartheta_G - \alpha\}\}$ for some α ; therefore if we substitute in the previous resolution $\overline{S}$ with T we get a graded minimal free resolution of a Gorenstein Artinian graded algebra A_G with $\vartheta_G = \vartheta_\Gamma$ (see [RZ1] Remark 3.8). Hence condition 2) is verified. Now observe that $\beta_G \leq \beta_\Gamma$ hence $\text{mci } \beta_G \leq \text{mci } \beta_\Gamma$ (see [RZ3] Theorem 3.9). Since $I_Z \subseteq I_\Gamma$ ($d_1, d_2, d_3 \geq \text{mci } \beta_\Gamma \geq \text{mci } \beta_G = (e_1, e_2, e_3)$). Let $s \in \text{Supp}(S \setminus T)$ and $i = \min\{j \mid d_j = s\} + \mu_{S \setminus T}(s) - 1$, say $m := \mu_{(S \setminus T)}(s)$. Then, among f_1, f_2, f_3 , there are m forms, of degree s , such that each of them is either a generator not minimal for I_Γ or it is a minimal generator for I_Γ , but there exists a minimal generator for I_Γ of degree $\vartheta_\Gamma - s$. So, by Lemma 3.4 and Lemma 3.5, $|\{j \mid d_j = s, d_j > e_j\}| \geq m$. Let $h := \max\{j \mid d_j = s, d_j > e_j\}$. Then

$$h \geq \min\{j \mid d_j = s, d_j > e_j\} + m - 1 \geq i := \min\{j \mid d_j = s\} + m - 1,$$

therefore $s = d_i = d_h > e_h \geq e_i$ and we are done.

Vice versa, since (D, E, F) is *aci*-type we have

- 1) $D = \{d_0\} \sqcup \overline{D} \sqcup S$, $|\overline{D} \sqcup S| = 3$;
- 2) $E = (d - F) \sqcup (d_0 + \overline{D}) \sqcup (\vartheta_Z - S)$.

Let $A_G = R/I_G$ be a 3-codimensional Artinian Gorenstein algebra with Betti sequence $\beta_G = (G_0, G_1, G_2)$; by condition 3), using part 2) of Lemma 3.4 for every $s \in \text{Supp}(S \setminus T)$ with multiplicity $\mu_{S \setminus T}(s) = m$ we can find a regular sequence of length 3 in I_G with m elements of degree s which are not minimal generators for I_G (and the other elements minimal generators for I_G). Say I_Z the complete intersection generated by this regular sequence and $I_Q := I_Z : I_G$; by mapping cone procedure we see that the minimal resolution of I_Q will be of the following type:

$$\begin{array}{c}
 R(-d_0) \oplus \bigoplus_{h \in \overline{D}} R(-h) \oplus \bigoplus_{h \in S} R(-h) \\
 \uparrow \\
 \bigoplus_{h \in d - F} R(-h) \oplus \bigoplus_{h \in d_0 + \overline{D}} R(-h) \oplus \bigoplus_{h \in \vartheta_Z - S} R(-h) \\
 \uparrow \\
 \bigoplus_{h \in F} R(-h) \\
 \uparrow \\
 0
 \end{array}$$

which means that (D, E, F) is a Betti sequence of an almost complete intersection. $\square$

Example 3.7. In this example we produce a sequence in which all the conditions of Theorem 3.6 hold but the last one, so it is not admissible as a Betti sequence for an almost complete intersection. Let $D = \{\{3, 6, 6, 6\}\}$, $E = \{\{8, 8, 8, 10, 10, 10, 12, 12, 12, 12\}\}$, $F = \{\{9, 11, 11, 11, 13, 13, 13\}\}$ and $\beta = (D, E, F)$. Then $\vartheta_Z = 18$, $\vartheta_G = 15$, $d = 21$, $D^* = \{\{6, 6, 6\}\}$, $S = \{\{6, 6, 6\}\}$, $G_0 = \{\{5, 5, 5, 7, 7, 7, 9\}\}$, $G_1 = \{\{6, 8, 8, 8, 10, 10, 10\}\}$, $G_2 = \{15\}$. Note that $\beta_G := (G_0, G_1, G_2)$ is a Gorenstein Betti sequence with $\text{mci } \beta_G = (5, 5, 7)$ hence $(6, 6, 6) \not\geq \text{mci } \beta_G$.

Example 3.8. In this example we produce a sequence in which all the conditions of Theorem 3.6 hold, except that one regarding the $i = \min\{j \mid d_j = s\} + \mu_{S \setminus T}(s) - 1$ for some $s \in \text{Supp}(S \setminus T)$, so it is not admissible as a Betti sequence for an almost complete intersection. Let $D = \{\{2, 5, 5, 7\}\}$, $E = \{\{7, 7, 7, 9, 9, 9, 10, 11\}\}$, $F = \{\{8, 10, 10, 10, 12\}\}$ and $\beta = (D, E, F)$. Then $\vartheta_Z = 17$, $\vartheta_G = 15$, $d = 19$, $D^* = \{\{5, 5, 7\}\}$, $S = \{7\}$, $G_0 = \{\{5, 5, 5, 7, 7, 7, 9\}\}$, $G_1 = \{\{6, 8, 8, 8, 10, 10, 10\}\}$, $G_2 = \{15\}$. Note that $\beta_G := (G_0, G_1, G_2)$ is a Gorenstein Betti sequence with $\text{mci } \beta_G = (5, 5, 7)$, $(d_1, d_2, d_3) = (5, 5, 7) \geq \text{mci } \beta_G$, but $3 = \min\{j \mid d_j = 7\} + \mu_{S \setminus T}(7) - 1$ and $d_3 = 7$ is not greater than $e_3 = 7$.

Example 3.9. The following sequences are admissible as Betti sequences for an almost complete intersection:

$$(D, E, F) = (\{4, 5^{(2)}, 9\}, \{9^{(3)}, 11^{(3)}, 13\}, \{12^{(3)}, 14\})$$

and

$$(D, E, F) = (\{4, 5^{(2)}, 9\}, \{9^{(3)}, 10, 11^{(3)}, 13\}, \{10, 12^{(3)}, 14\}).$$

Note that we can obtain the first one by the second one by deleting the “ghost” degree 10.

REFERENCES

- [BE] D. A. Buchsbaum, D. Eisenbud, *Algebra structures for finite free resolutions, and some structure theorems for ideals of codimension 3*, Amer. J. Math. **99**(1) (1977), 447–485.
- [Di] S. Diesel, *Irreducibility and dimension theorems for families of height 3 Gorenstein algebras*, Pac. J. of Math. **172**(4) (1996), 365–397.
- [Ga] F. Gaeta, *Quelques progrès récents dans la classification des variétés algébriques d’un espace projectif*, Deuxième Colloque de Géométrie Algébrique, Liège, (1952), pp. 145–183.
- [IK] A. Iarrobino, V. Kanev, *Power Sums, Gorenstein Algebras, and Determinantal Loci*, LNM 1721 (1999) Springer-Verlag
- [Mi] J. Migliore, *Introduction to liaison theory and deficiency modules*, Progress in Math. v.165 Birkhauser Boston (1998)
- [MM] J. Migliore, R. Mirò Roig, *On the minimal free resolution of $n + 1$ general forms*, Trans. Am. Math. Soc., 355 (2003) vol.1, 1–36
- [PS] C. Peskine, L. Szpiro, *Liaison des variétés algébriques. I*, Inv. Math. **26** (1974), 271–302.
- [RZ1] A. Ragusa, G. Zappalà, *Properties of 3-codimensional Gorenstein schemes*, Comm. Algebra, 29 (1), 303–318 (2001)
- [RZ2] A. Ragusa, G. Zappalà, *Gorenstein schemes on general surfaces*, Nagoya Math. J. Vol.162, 111–125 (2001)

- [RZ3] A. Ragusa, G. Zappalà, *Complete intersections containing Cohen Macaulay and Gorenstein schemes* Preprint.
- [Se] S. Seo *Almost complete intersections*, J. of Algebra **320** (2008), 2594–2609.

(A. RAGUSA) DIP. DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI CATANIA,
VIALE A. DORIA 6, 95125 CATANIA, ITALY
E-mail address: `ragusa@dmf.unict.it`
Fax number: +39095330094

(G. ZAPPALÀ) DIP. DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI CATANIA,
VIALE A. DORIA 6, 95125 CATANIA, ITALY
E-mail address: `zappalag@dmf.unict.it`
Fax number: +39095330094